

A NEW VIEW OF NOVAE: RADIO OBSERVATIONS OF V1723 AQL


JENNIFER WESTON
COLUMBIA UNIVERSITY

WITH THE E-NOVA PROJECT:
JENO SOKOLOSKI, LAURA CHOMIUK, THOMAS NELSON, MICHAEL RUPEN,
MIRIAM KRAUSS, AMY MIODUSZEWSKI, KOJI MUKAI, NIRUPAM ROY,
STEWART EYRES, MIKE BODE, TIM O'BRIEN, YONG ZHENG


V1723 AQL

- Went into outburst September 11th, 2010
- Optical implies a typical ‘fast’ nova
- $v \sim 1500$ km/s
- Neither an embedded nor a recurrent nova
- First northern classical nova after expansion of VLA bright enough for detailed observation


RADIO EMISSION FROM NOVAE



RADIO EMISSION FROM NOVAE




RADIO EMISSION FROM NOVAE










RADIO EMISSION FROM NOVAE

- Thermal Free-Free Emission
- Spherical Shell
- smooth density $\propto 1/r^2$
- Constant $T_e \sim 10^4$ K
- distribution of velocities




V1723 Aql







V1723 Aql




V1723 Aql




Difference between observations and thermal-only shell




Difference between observations and thermal-only shell





V1723 Aql



J2000 Declination

